

09 013329

EPO - DG 1

518 Rec'd PCT/PTO 13 AUG 2001 13. 04. 2001

WHAT IS CLAIMED IS:

(82)

1. A method for obtaining a Neprilysin-like (NEP-like) metallopeptidase which comprises the following steps:
- 5 – selecting a primer in C-terminus of the His-Glu-Xaa-Xaa-His (where Xaa represents any amino acid) with a degenerate nucleotide sequence complementary to at least the Gly-Glu-Asn-Ile-Ala-Asp amino acid sequence of known NEP-like metallopeptidases with sufficient binding capacity;
 - 10 – selecting a primer in N-terminus of the His-Glu-Xaa-Xaa-His (where Xaa represents any amino acid) with a degenerate nucleotide sequence complementary to a conserved amino acid sequence with preferably 80% homology with known NEP-like metallopeptidases and sufficient binding capacity;
 - 15 – contacting said primer with tissue nucleic acids to yield PCR products;
 - selecting said PCR products that contain the His-Glu-Xaa-Xaa-His motif; and
 - completing the sequence of said selected PCR products with standard methods.
- 20 2. A metallopeptidase sharing about 80% homology with the amino acid sequence shown in Figure 3.
3. A metallopeptidase which is soluble sharing about 80% homology with the amino acid sequence in C-terminus of the furin site shown in Figure 3.
4. A metallopeptidase which is soluble sharing about 80% homology with the amino acid sequence shown in Figure 3 and with an enzymatic activity capable of degradation of known Neprilysin substrates, preferably
25 Tyrosyl-[3,5-³H1](D-Ala₂)-Leu₅-enkephalin and bradykinin.
5. A composition comprising a metallopeptidase as defined in any one of claims 2-4.

FOR INFO SEE FIG 3

SW
RT



6. A nucleic acid encoding a metallopeptidase as defined in any one of claims ~~2-4~~.

7. An antibody directed against a metallopeptidase as defined in any one of claims ~~2-4~~.

5 8. A method for obtaining a substrate of a metallopeptidase as defined in any one of claims ~~2-4~~, which comprises the steps of:

-- contacting said metallopeptidase with a molecule or extract; and

-- assaying the resulting solution for a decrease in said molecule or extract, when compared with the same but in absence of said metallopeptidase, as an indication of the presence of said substrate.

10

9. A method for obtaining an inhibitor of a metallopeptidase as defined in any one of claims ~~2-4~~, which comprises the steps of:

-- contacting said metallopeptidase with a molecule or extract in the presence of a substrate selected known NEP substrates, preferably Tyrosyl-[3,5-³H1](D-Ala₂)-Leu₅-enkephalin and bradykinin; and

15

-- assaying the resulting solution for an increase in said substrate, when compared with the same but in absence of said metallopeptidase, as an indication of the presence of said inhibitor.

Handwritten note: "Note" with a circle around the number 10 and an asterisk.

10. An inhibitor obtained from the method of claim 9.

20

11. The use of a known NEP inhibitor or an inhibitor as defined in claim 10 to control the enzymatic activity of a metallopeptidase as defined in any one of claims ~~2-4~~.

Handwritten note: "SW2 A2" with a bracket pointing to claim 11.

12. The use of a known NEP inhibitor or an inhibitor as defined in claim 10 to manage disease relating to the physiological status of the cardiovascular system, the central nervous system, the spleen, the liver, the kidney, the male reproductive system or the maturation of spermatozoa.

25

13. The use of a metallopeptidase as defined in any one of claims ~~2-4~~ to manage disease relating to the physiological status of the cardiovascular

Handwritten note: "SW3 A3" with a bracket pointing to claim 13.

FOI b7D - 6222660

system, the central nervous system, the spleen, the liver, the kidney, the male reproductive system or the maturation of spermatozoa.

14. A recombinant vector comprising 80% homology with the nucleic acid encoding the N-terminal part of the amino acid sequence shown in Figure 3, which N-terminal part terminates with a furin-recognition sequence.

5

15. A method for producing a soluble form of a protein, polypeptide or part thereof which comprises:

- obtaining nucleic acids encoding said protein, polypeptide or part thereof;

10

- fusing said nucleic acids in phase with an N-terminal fragment wherein said N-terminal fragment comprises a cleavable furin-like site located in C-terminus past the transmembrane region or is an N-terminal part as defined in claim 14;

15

- having the fused nucleic acids to be expressed in a host cell which expresses or is made to express furin in the presence of a culture medium; and

- recovering said soluble form in the culture medium.

20

16. A protein, polypeptide or part thereof produced by the method defined in claim 15, wherein said protein, polypeptide or part thereof is a metallopeptidase sharing about 80% homology with the region in C-terminus of the putative furin site of the amino acid sequence shown in Figure 4.

25

17. A metallopeptidase sharing about 80% homology with the region in C-terminus of the putative furin site of the amino acid sequence shown in Figure 4.

18. A metallopeptidase sharing about 80% homology with the region in C-terminus of the putative furin site of the amino acid sequence shown in Figure 4 and with an enzymatic activity capable of degradation of known

NEP substrates, preferably Tyrosyl-[3,5-³H1](D-Ala₂)-Leu₅-enkephalin and bradykinin.

19. A composition comprising a metallopeptidase as defined in any one of claims 16-18.

5 20. A nucleic acid encoding a metallopeptidase as defined in any one of claims 16-18.

21. An antibody directed against a metallopeptidase as defined in any one of claims 16-18.

SWS
AS

10 22. A method for obtaining a substrate of a metallopeptidase as defined in any one of claims 16-18, which comprises the steps of:

- contacting said metallopeptidase with a molecule or extract; and
- assaying the resulting solution for a decrease in said molecule or extract, when compared with the same but in absence of said metallopeptidase, as an indication of the presence of said substrate.

15 23. A method for obtaining an inhibitor of a metallopeptidase as defined in any one of claims 16-18, which comprises the steps of:

- contacting said metallopeptidase with a molecule or extract in the presence of a substrate selected from known NEP substrates or a protein, polypeptide or part thereof produced by the method of claim 15, preferably Tyrosyl-[3,5-³H1](D-Ala₂)-Leu₅-enkephalin and bradykinin; and
- assaying the resulting solution for an increase in said substrate, when compared with the same but in absence of said metallopeptidase, as an indication of the presence of said inhibitor.

20

24. An inhibitor obtained from the method of claim 23.

25 25. The use of a known NEP inhibitor or an inhibitor as defined in claim 24 to control the enzymatic activity of a metallopeptidase as defined in any one of claims 16-18.

SWS
AS

FOOTNOTES

SW
9/6

43

FORERO SEE 1660

- 26. The use of a known NEP inhibitor or an inhibitor as defined in claim 24 to manage disease relating to the physiological status of the cardiovascular system, the central nervous system, the spleen, the liver, the kidney, the male reproductive system or the maturation of spermatozoa.
- 5 27. The use of a metallopeptidase as defined in any one claims ~~16-18~~ to manage disease relating to the physiological status of the cardiovascular system, the central nervous system, the spleen, the liver, the kidney, the male reproductive system or the maturation of spermatozoa.
- 10 28. A method as defined in claim 15, wherein said protein, polypeptide or part thereof is beta-endorphin.
- 15 29. A recombinant host cell capable of expressing a protein, polypeptide or part thereof transplanted in a mammal to manage a disease, physiological process or pain.
- 30. A metallopeptidase sharing about 80% homology with the amino acid sequence located in the C-terminus of the predicted transmembrane domain of the amino acid sequence shown in Figure 5 which has been produced by the method of claim 15, by fusing in frame a cleavable signal peptide in N-terminus of said amino acid sequence or by transforming said predicted transmembrane domain into a cleavable signal peptide.
- 20 31. A composition comprising a metallopeptidase as defined in claim 30.
- 32. An antibody directed against a metallopeptidase as defined in claim 30.
- 33. A method for obtaining a substrate of a metallopeptidase as defined in claim 30, which metallopeptidase shares about 80% homology with the C-terminal region of the predicted transmembrane domain of the amino acid sequence shown in Figure 5, comprising the steps of:
 - contacting said metallopeptidase with a molecule or extract; and
- 25



-- assaying the resulting solution for a decrease in said molecule or extract, when compared with the same but in absence of said metallopeptidase, as an indication of the presence of said substrate.

5 34. A method for obtaining an inhibitor of a metallopeptidase sharing about 80% homology with the C-terminal region of the predicted transmembrane domain of the amino acid sequence shown in Figure 5, which comprises the steps of:

-- contacting said metallopeptidase with a molecule or extract in the presence of a substrate produced by the method of claim 33; and

10 -- assaying the resulting solution for an increase in said substrate, when compared with the same but in absence of said metallopeptidase, as an indication of the presence of said inhibitor.

35. An inhibitor obtained by the method of claim 34.

15 36. The use of an inhibitor as defined in claim 35 to control the enzymatic activity of the metallopeptidase sharing about 80% homology with the C-terminal region of the predicted transmembrane domain of the amino acid sequence shown in Figure 5.

20 37. The use of an inhibitor as defined in claim 35 to manage disease relating to the physiological status of the central nervous system, the spleen or the bones.

38. The use of a metallopeptidase as defined in claim 30 to manage disease relating to the physiological status of the cardiovascular system, the central nervous system, the spleen or the bones.

FOR 230 SEE 1650

Add Art 7